

Two-dimensional electron gas in uniform magnetic and electric fields

Choon-Lin Ho¹, V. R. Khalilov² and Chi Yang¹

¹Department of Physics, Tamkang University, Tamsui, Taiwan 25137, R.O.C.

² Department of Theoretical Physics, Physical Faculty,
Moscow State University, 119899, Moscow, Russia.

Abstract

The thermodynamic potential of an ideal nonrelativistic gas of two-dimensional electrons in crossed uniform magnetic and electric fields is constructed. For low temperatures and very weak electric fields, it is shown that the Hall conductance is always quantized at integral multiples of e^2/h over a large range of strong magnetic fields. This could be viewed as a quantum statistical explanation of the integral quantum Hall effect. Magnetic properties of the system at high temperatures are briefly discussed.

PACS. — 73.40.Hm, 71.10.Ca, 05.30.-d

Two-dimensional electron gas (2DEG) in external electromagnetic fields at low temperature has attracted significant interests since the discovery of the integral quantum Hall effect (IQHE) [1]. In IQHE, the Hall conductance σ_{yx} was discovered to have a stepwise dependence (the appearance of plateaux) on the strong external magnetic field. At these plateaux, σ_{yx} is quantized to $\sigma_{yx} = ne^2/h$, $n = 1, 2, \dots$ (e is the magnitude of the electron charge, and $h = 2\pi\hbar$ is the Planck constant) to an accuracy as high as several parts in 10^8 , while the longitudinal conductivity σ_{xx} nearly vanishes. The vanishing of σ_{xx} implies the absence of dissipation. The amazing fact that the Hall conductance is quantized to such a high precision, and that the quantized values are related only to fundamental constants and are independent of the actual sample and geometry, indicate that IQHE must be explicable from some general principles independent of any finer details of the interaction mechanism in the sample. Efforts in this direction in the past have resulted in theory of IQHE based on the principle of gauge invariance [2] and on topological considerations [3].

In view of the fact that IQHE is independent of interaction, and that it is a lossless effect as implied by the vanishing of σ_{xx} , one may wonder if IQHE could also be deducible from the quantum statistics of a non-interacting 2DEG at thermodynamic equilibrium in the crossed magnetic and electric fields. The purpose of this paper is to investigate such possibility.

Let us consider a system of nonrelativistic non-interacting 2DEG in a crossed uniform electric ($\mathbf{E}$) and magnetic ($\mathbf{B}$) fields, which are specified in Cartesian coordinates as

$$\mathbf{E} = (0, E, 0), \quad \mathbf{B} = (0, 0, B) = \nabla \times \mathbf{A}, \quad \mathbf{A} = (-yB, 0, 0). \quad (1)$$

We shall consider only the case of very weak electric field and strong magnetic field ($E \ll B$).

The energy spectrum of two-dimensional electron in the field configuration (1) is given by (see e.g. [4, 5]):

$$E_n = \frac{\hbar e B}{mc} \left(n + \frac{1}{2} \right) - c \frac{E}{B} \tilde{p} + m_s g \frac{\hbar e B}{2m_0 c}, \quad m_s = \pm \frac{1}{2}, \quad (2)$$

where n is the Landau level number, m is the effective electron mass in a crystal lattice, m_0 is the mass of a free electron, g is the effective Landé factor, and $\tilde{p}$ is the momentum of electron in the x -direction. For a very weak electric field, $\tilde{p}$ is constrained by $|\tilde{p}| \leq eBL_y/2c$ [4]. The last term in (2) describes the interaction of the spin magnetic moment of the electron with the magnetic field. If the electrons are free we shall put $m = m_0$ and $g = 2$.

The main physical quantity which defines all thermodynamical properties of noninteracting particles gases is the thermodynamic potential (TP) $\Omega(\tilde{\mu}, B, E, T)$, considered as a function of the chemical potential ($\tilde{\mu}$), the external fields and the gas temperature (T). The TP of the 2DEG in the XOY plane can be defined as follows (see e.g.: [6]):

$$\Omega(\tilde{\mu}, B, E, T) = -k_B T \sum_{n, \tilde{p}, m_s} \ln \left[1 + \exp \left(\frac{\tilde{\mu} - E_n}{k_B T} \right) \right], \quad (3)$$

where k_B is the Boltzmann constant. In terms of dimensionless quantities $p \equiv \tilde{p}/mc$, $\mu \equiv \tilde{\mu}/mc^2$, $\varepsilon_n \equiv E_n/mc^2$ and $\theta \equiv k_B T/mc^2$, Ω is given by

$$\Omega(\mu, B, E, \theta) = -mc^2 n^{(2)} \theta \int_{-b/2}^{b/2} dp \frac{1}{b} \sum_{n, m_s} \ln \left[1 + \exp \left(\frac{\mu - \varepsilon_n}{\theta} \right) \right]. \quad (4)$$

Here L_x , L_y are the lengths of the two-dimensional space, $b = eBL_y/mc^2$, and

$$n^{(2)} = \frac{eBL_x L_y}{hc} \quad (5)$$

is the number of quantum electron states in the magnetic field for a given n in an area $L_x L_y$.

Evaluation of the TP is easily carried out by the Mellin transformation method with respect to the variable $\exp(\mu/\theta)$, resulting in the following expression

$$\begin{aligned} \Omega(\mu, \kappa, E, \theta) = & \mp mc^2 \theta n^{(2)} \sum \text{Res} \frac{\pi \exp(s\mu/\theta)}{s \sin(\pi s)} \\ & \times \frac{\cosh[sg^* \kappa/2\theta]}{\sinh(s\kappa/2\theta)} \frac{\sinh[seEL_y/2mc^2\theta]}{seEL_y/2mc^2\theta}, \\ & g^* = \frac{gm}{2m_0}, \quad \kappa = \frac{B}{B^*}, \quad B^* = \frac{m^2 c^3}{e\hbar}, \end{aligned} \quad (6)$$

where the upper (lower) sign refers to closing the contour to the left (right) of the imaginary axis for $\mu > 0$ ($\mu < 0$). When $\mu > 0$, the sum of residues of (6) should be calculated at the poles on both the negative real semiaxis $s = 0, -1, -2, -3, \dots$ (including $s = 0$) and the imaginary axis $s_l = 2\pi il\theta/\kappa$. For $\mu < 0$, the integration contour is closed to the right and the sum of residues should be calculated at the poles $s = 1, 2, 3, \dots$

The poles located on the real axis contribute to the “monotonic part” of Ω , while the poles $s_l = 2\pi il\theta/\kappa$ dominate the behavior of its oscillating part. Since we are interested in an almost degenerate 2DEG, (6) must be calculated in the limit $\theta \ll \mu$, in which case $\mu > 0$. It is important that values of the chemical potentials μ may exceed zero only for fermions. It follows that only thermodynamic quantities characterizing fermions in a magnetic field at low temperature may exhibit the oscillating behavior (the so-called Landau oscillations).

Let us first consider the case of low temperature limit when $\theta \ll \kappa$. Taking the sum at the poles $s_l = 2\pi il\theta/\kappa$, and the pole $s = 0$, we obtain TP to the first order in θ/κ in the form

$$\begin{aligned}\Omega &= \Omega_{\text{mon}} + \Omega_{\text{osc}} \quad , \\ \Omega_{\text{mon}} &= -mc^2 n^{(2)} \kappa \left[\frac{z^2}{4\pi^2} + \frac{1}{4} \left(g^{*2} - \frac{1}{3} \right) - \frac{\gamma\theta}{\pi\kappa} z + \frac{\alpha^2}{12} \right] \quad , \\ \Omega_{\text{osc}} &= -mc^2 n^{(2)} \kappa \sum_{l=1}^{\infty} (-1)^{l+1} \frac{\cos(g^* \pi l) \cos(zl) \sin(\alpha \pi l)}{\pi^2 l^2 \alpha \pi l} \quad ,\end{aligned}\tag{7}$$

where $z = 2\pi\mu/\kappa$, $\alpha = eEL_y/\kappa mc^2$ and $\gamma = 0.57721$ is the Euler constant. We note here that in deriving the monotonic part of TP (Ω_{mon}), we took into account only the contribution of the pole $s = 0$, since the poles $s \neq 0$ contribute to terms of the order $(-1)^{l+1} \exp \{ - [l(\mu - (g^* - 1)\kappa/2) / \theta] \} / l$, which are exponentially small.

In the case of very weak electric fields such that $\alpha \ll 1$, *i.e.* for E satisfying

$$eEL_y \ll \kappa mc^2 \quad ,\tag{8}$$

the TP in (7) reduces to

$$\begin{aligned}\Omega &= -mc^2 n^{(2)} \kappa \left[\frac{z^2}{4\pi^2} + \frac{1}{4} \left(g^{*2} - \frac{1}{3} \right) - \frac{\gamma\theta}{\pi\kappa} z + \frac{1}{\pi^2} F(z) \right], \\ F(z) &= \sum_{l=1}^{\infty} (-1)^{l+1} \frac{1}{l^2} \cos(g^* \pi l) \cos(zl) .\end{aligned}\quad (9)$$

Having derived the TP of the electron gas, we can then calculate the number of particles

$$N_e = -\frac{\partial\Omega}{mc^2\partial\mu}, \quad (10)$$

and the magnetic moment

$$\mathbf{M} = -\frac{\partial\Omega}{\partial\mathbf{B}} \quad (11)$$

of the electron gas in fields (1) at thermodynamic equilibrium. From (9) and (11) the magnetic moment of the system is

$$\begin{aligned}M_x &= M_y = 0 \\ M_z &= mc^2 \mu \frac{eL_x L_y}{\hbar c} \left[\frac{1}{2} \left(g^{*2} - \frac{1}{3} \right) \frac{1}{z} + \frac{2F(z)}{z\pi^2} - \frac{1}{\pi^2} \frac{dF(z)}{dz} \right].\end{aligned}\quad (12)$$

One sees that the oscillating part of the magnetic moment dominate in the limit $\mu \gg \kappa$. This is the Landau-de Haas-van Alphen effect [6].

Now we shall evaluate the number of electrons N_e according to (10) with the TP given by (9). This quantity is of great importance for the problem under consideration. In fact, the off-diagonal Hall conductivity in fields (1) can be related to the number of two-dimensional electrons N_e that occupy quantum states with $\varepsilon_n < \varepsilon_F$ (where $\varepsilon_F = \tilde{\mu}$ at $\theta = 0$ is the Fermi level) at thermodynamic equilibrium as follows:

$$\sigma_{yx} = \frac{ecn_e}{B} \equiv \frac{e^2}{h} \cdot \frac{N_e}{n^{(2)}}, \quad (13)$$

where $n_e = N_e/L_x L_y$ is the density of electrons. The ratio

$$\Phi(z) \equiv \frac{N_e(z)}{n^{(2)}} \quad (14)$$

as a function of z characterizes the number of occupied Landau levels, and is usually called the filling factor. From (9) and (10), we get (for $\theta \ll \kappa$):

$$\Phi(z) = \frac{1}{\pi} \left(z + 2 \frac{dF(z)}{dz} \right) . \quad (15)$$

The functions $F(z)$ in (9) is a piecewise periodic function of z . To see this, we express the product of cosines in (9) as a sum of cosines, and use the identity [7]:

$$\sum_{l=1}^{\infty} (-1)^{l+1} \frac{\cos(lx)}{l^2} = \frac{\pi^2}{12} - \frac{x^2}{4} , \quad -\pi \leq x \leq \pi . \quad (16)$$

The result is:

$$F(z) = \begin{cases} \frac{\pi^2}{12} - \frac{1}{4}z^2 - \frac{1}{4}g^{*2}\pi^2 , & -(1-g^*)\pi \leq z \leq (1-g^*)\pi ; \\ \frac{\pi^2}{12} + \frac{\pi}{2}z - \frac{1}{4}z^2 - \frac{1}{4}g^{*2}\pi^2 - \frac{1}{2}(1-g^*)\pi^2 , & (1-g^*)\pi \leq z \leq (1+g^*)\pi . \end{cases} \quad (17)$$

Substituting (17) into (15), one finds:

$$\Phi(z) = \begin{cases} 0 , & 0 \leq z \leq (1-g^*)\pi ; \\ i , & (i-g^*)\pi \leq z \leq (i+g^*)\pi ; \\ i+1 , & (i+g^*)\pi \leq z \leq (i+2-g^*)\pi ; \end{cases} \quad (18)$$

$$i = \text{odd integers} = 1, 3, 5, \dots$$

Hence the filling factor Φ assumes only positive integral values. Consequently, from (13), we see that the Hall conductance σ_{yx} is quantized at integral multiples of e^2/h . IQHE can thus be explained from the quantum statistical nature of the 2DEG. The widths of the plateaux corresponding to odd filling factors Φ are equal to $2g^*\pi$, while those corresponding to even Φ are equal to $2(1-g^*)\pi$. Two special cases, namely, $g^* = 0$ and $g^* = 1$, are worthy of note. From (18) it is clear that when $g^* = 0$ the filling factor Φ is always an even integer, and when $g^* = 1$, Φ is always odd. For $g^* = 0$ there is no splitting of spin degree of freedom in each Landau level, and the σ_{yx} 's are even integral multiples of e^2/h [8]. For $g^* = 1$, all Landau levels except the ground state are spin degenerate, and σ_{yx} 's are odd integral

multiples of e^2/h . In actual samples exhibiting IQHE, g^* are usually much smaller than unity. For instance, $m \approx 0.07m_0$ and $g \approx 0.8$ in GaAs, giving $g^* \approx 0.03$ [9]. Thus one expects from (18) that for such small value of g^* , the odd plateaux are much narrower than the even ones. This explains why the experimentally observed plateaux mostly correspond to even integers [10].

One could also obtain the quantized values of σ_{yx} in layered multi-quantum-well structures. These structures contain a number of well separated identical two-dimensional electronic systems. If the number of layers in such a structure is J , then the total number of electrons is $N_e^T = JN_e$, where N_e is the number of electrons in a single layer given by $N_e = n^{(2)}\Phi = n^{(2)}I$, $I = 1, 2, 3, \dots$, according to (14) and (18). Hence the Hall conductance of the system is

$$\sigma_{yx} = \frac{e^2}{h} \cdot \frac{N_e^T}{n^{(2)}} = \frac{e^2}{h} IJ, \quad (19)$$

which agreed with the experimental values [11].

Let us now consider briefly the behaviour of 2DEG at high temperature ($\theta \gg \kappa$) for a very weak electric field satisfying the condition (8), and for $E = 0$. In these cases, the TP is easily obtained from eq. (6) by closing the contour to the right of the imaginary axis and taking the lower sign. The residues are evaluated at the poles $s = 1, 2, \dots$. The result is:

$$\Omega = mc^2\theta n^{(2)} \sum_{l=1}^{\infty} (-1)^l \frac{\exp(l\mu/\theta)}{l} \frac{\cosh(lg^*\kappa/2\theta)}{\sinh(l\kappa/2\theta)}. \quad (20)$$

We can also obtain, by the same method described above, the TP of a two-dimensional nonrelativistic spin-0 free boson gas at arbitrary temperature. We give the result here for reference (see also [12, 13])

$$\Omega_0 = -\frac{1}{2}mc^2\theta n^{(2)} \sum_{l=1}^{\infty} \frac{\exp(l\mu/\theta)}{l \sinh(l\kappa/2\theta)}. \quad (21)$$

The factor of $1/2$ accounts for the difference in the spin degree of freedom. We note here that all the chemical potentials in (20) and (21) are assumed to be less than zero. It is

worth emphasizing that (21) is valid at low temperature also, since for bosons one must always close the contour to the right of the imaginary axis. From (21), (10) and (11) we can obtain the boson number and magnetic moment density, and reproduce the results of [12, 13] concerning two-dimensional ideal nonrelativistic spin-0 boson gas at low temperature.

At very high temperature equations (20) and (21) reduce, up to the first order term in κ/θ , to

$$\Omega \cong mc^2\theta n^{(2)} \left[\frac{2\theta}{\kappa} \text{Li}_2(-e^{\mu/\theta}) - \frac{1}{4} \left(g^{*2} - \frac{1}{3} \right) \frac{\kappa}{\theta} \frac{1}{\exp(-\mu/\theta) + 1} \right], \quad (22)$$

$$\Omega_0 \cong -\frac{1}{2}mc^2\theta n^{(2)} \left[\frac{2\theta}{\kappa} \text{Li}_2(e^{\mu/\theta}) - \frac{1}{12} \frac{\kappa}{\theta} \frac{1}{\exp(-\mu/\theta) - 1} \right]. \quad (23)$$

Here $\text{Li}_2(x) = \sum_{l=1}^{\infty} x^l/l^2$ is the dilogarithmic function of x .

We see that the first terms on the right hand sides of (22) and (23) contribute only in the densities of the particles, while the second terms give contribution both in the densities of the particles and the densities of the magnetic moments of the gases.

From (22), (23) and (11) one can check that the spin-0 and spin-1/2 (with $g^* < 1/\sqrt{3}$) particles gases produce magnetic moments that direct against the external magnetic field induction. They are thus diamagnetic in nature. When $g^* > 1/\sqrt{3}$, the electron gas is paramagnetic.

In summary, we have shown that the two main features of IQHE, namely the appearance of a sequence of plateaux in the Hall conductivity σ_{xy} , and the exact quantization of σ_{xy} at integral multiples of e^2/h , can be deduced directly in the framework of quantum statistics from an equilibrium non-interacting 2DEG in crossed magnetic and electric field in low temperature. At high temperature, there exists a critical value of $g_{cr}^* = 1/3$ below which the system is diamagnetic. Though we have assumed in our derivation a rectangular two-dimensional space, the results are independent of the geometry of the space in the thermodynamic limit.

Acknowledgment

This work is supported in part by the R.O.C. Grants number NSC-85-2112-M-032-002.

References

- [1] R.E. Prange and S.M. Girvin (eds.), *The Quantum Hall Effect*, 2nd ed., Springer-Verlag, New York, 1990.
- [2] R.B. Laughlin, *Phys. Rev.* **B23**, 5632 (1981).
- [3] D. Thouless, in ref. [1].
- [4] L.D.Landau and E.M.Lifshitz, *Quantum Mechanics*, 3rd ed., Pergamon, London, 1977.
- [5] V.R. Khalilov, *Electrons in Strong Electromagnetic Fields: an advanced classical and quantum treatment*, Gordon & Breach Sci. Pub., Amsterdam, 1996.
- [6] L.D. Landau and E.M.Lifshitz, *Statistical Physics*, Part I, 3rd ed., Pergamon, London, 1980.
- [7] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, 5th ed., Academic Press, San Diego, 1994.
- [8] S. Datta, *Electronic Transport in Mesoscopic Systems*, Cambridge University Press, Cambridge, 1995.
- [9] H.L. Störmer, in *Frontiers in Physics, High Technology and Mathematics*, (Eds.) H.A. Cerdeira and S.O. Lundqvist, World Scientific, Singapore, 1990.
- [10] M.E. Cage, R.F. Dziuba and B.F. Field, *IEEE Trans. Instrum. Meas.* **IM-34**, 301 (1985).
- [11] T. Haavasoja, H.L. Störmer, D.J. Bishop, V. Narayanamurti, A.C. Gossard, and W. Wiegmann, *Surf. Sci.*, **142**, 294 (1984).
- [12] R.M. May, *Phys. Rev.* **115**, 254 (1959).

- [13] D.J. Toms, *Phys. Rev.* **D51**, 1886 (1995).